

# AIRS v6 Level-2 Startup State

Evan Manning  
George Aumann  
Jan Gohlke

Jet Propulsion Laboratory  
California Institute of Technology

# What It Is



- The AIRS physical retrieval works on cloud-cleared radiances.
- These radiances come from a cloud-clearing step.
- The cloud-clearing step in turn needs to start with an estimate of atmospheric  $T(p)$ ,  $q(p)$ 
  - Use of IR in this stage raises issues of duplicate use of information content.

# History

- AIRS has always used a 3-step startup process:
  - ① A step which gives a rough  $T(p)$ ,  $q(p)$  using only MW and/or **cloudy** radiances
  - ② An initial cloud clearing
  - ③ A regression
- Up to v4 step #1 was a MW-only optimal estimation retrieval based on a NCEP/UARS  $T(p)$ ,  $q(p)$  climatology
- For v5 step #1 changed to a regression that works on cloudy radiances
  - An IR-only cloudy regression was originally developed for v5 IR-only retrieval.
  - Performance was so good that an IR+MW cloudy regression was developed for the IR+MW system.

# Why Revisit Startup Now?

- An improved ECMWF forecast based climatology is now available
- Some limitations of the current startup approach have emerged
  - Does not provide internal error estimates to later retrieval steps
  - Does not provide averaging kernels
  - Significant work is required for maintenance
  - Can react unpredictably to novel data:
    - Spectral shifts
    - Increasing CO<sub>2</sub>
  - Is a likely contributor to spurious trends in retrieved products
- Bill Blackwell of MIT has produced a new candidate startup algorithm: SCCNN
  - Stochastic cloud clearing
  - Neural network
  - Reported at previous science team meetings

# ECMWF-Based Climatology

- The make-up of the ECMWF forecast climatology was described in detail at the 2009-02 netmeeting  
[https://airsteam.jpl.nasa.gov/netmeeting/20090226.Netmeeting/Manning\\_ECMWF\\_AIRSv5\\_clim.pdf](https://airsteam.jpl.nasa.gov/netmeeting/20090226.Netmeeting/Manning_ECMWF_AIRSv5_clim.pdf)
- This climatology leverages existing programs which:
  - Extract ECMWF forecast/analysis at AIRS observation locations into AIRS-compatible 100-layer products
  - Grid these products in 1x1 degree cells

# Climatology Comparison

|                                | Old   | New  |
|--------------------------------|---|--|
| Spatial Resolution - Latitude  | 2.5 degrees (troposphere)<br>10 degrees (stratosphere)  | 1 degree                                   |
| Spatial Resolution - Longitude | 2.5 degrees (troposphere)<br>360 degrees (stratosphere) | 1 degree                                   |
| Temporal Resolution            | 1 month   | 1 month                                    |
| Temporal Specificity           | Variation within a day not modeled                      | Appropriate for local overpass time        |
| Surface type                   | No differentiation                                      | Land/Sea handled separately                |
| Model                          | NCEP 15-year reanalysis                                 | ECMWF forecast ~2008 with improved surface |
| Variance                       | No variance data  | Has variance data                          |

# Using the New Climatology

- The new climatology could be used in several ways:
  - Input to the MW-only retrieval in the V4 architecture
  - Replace the MW-only or cloudy regression as the first stage of startup
  - Replace the entire startup
- The exercise reported next used the climatology in place of the entire startup

# Exercise Description

- These runs use simulated data from UMBC for 2008-12-08
  - Surface is Masuda land & ocean
  - There are clear and cloudy radiances for the entire day
    - This is a high quality simulation
    - According to George Aumann the clouds look realistic
      - See his presentation on Wednesday
  - The truth is known
- The baseline algorithm (v5.4) is compared to the same algorithm with climatology startup replacing both regressions (v5.4Clim)
  - Both have Joel Susskind's surface improvements
  - Both are IR-only
- V5.4Clim has climatology naively “dropped in”
  - No use is made of the variance
  - No adjustments have been made to the physical retrieval
  - The climatology used for startup is based on December 2007
    - Using December 2008 would be cheating




# Exercise Description

- Because QC has not been adapted for v5.4Clim, all comparisons are performed over all retrievals.
  - Yield is 100%
  - This is a preliminary proof of concept, not a robust test
- Skill is used for the comparison.
  - $\text{Skill} = \text{correl}(\text{retrieved} - \text{background}, \text{truth} - \text{background}) * \sqrt{\text{yield}}$
  - See talk  
[http://airs.jpl.nasa.gov/documents/science\\_team\\_meeting\\_archive/science\\_team\\_meeting\\_2008.10.14/science\\_team\\_meeting\\_2008.10.14\\_files/mannning.2008-10-18.pdf](http://airs.jpl.nasa.gov/documents/science_team_meeting_archive/science_team_meeting_2008.10.14/science_team_meeting_2008.10.14_files/mannning.2008-10-18.pdf)
  - Skill effectively weights cases more highly the more they diverge from a climatology background
  - The same climatology which is used as a first guess in v5.4Clim is used as a background in the skill calculation.
    - This means that if the physical retrieval passed the climatology through unaltered it would get a skill of zero.

# Exercise Limitations


- The implementation of v5.4Clim was incomplete
- Only one day of data was used
- Only simulated data was used
- The lack of QC makes interpretation of results difficult.

# Clear Results: Ascending SST



- Results of v5.4 and v5.4Clim are comparable
  - Both have ~1 K warm bias
    - This may be due to details of the simulation (did not include optical depth tuning)
  - Both have trouble at high scan angles in very wet or dry cases
- v5.4 is noisier than v5.4Clim and has more angle dependence
- v5.4Clim has larger regions where T<sub>surf</sub> is biased significantly cool
  - These are dry regions relative to climatology






National Aeronautics and  
Space Administration  
  
Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

*Atmospheric Infrared Sounder*

v5.4

SST - Truth

# Clear Results: Ascending SST



5/4/09

Copyright 2009 California Institute of Technology.  
Government sponsorship acknowledged.

12






National Aeronautics and  
Space Administration  
Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

*Atmospheric Infrared Sounder*

v5.4 Clim

SST - Truth

# Clear Results: Ascending SST




5/4/09

Copyright 2009 California Institute of Technology.  
Government sponsorship acknowledged.

13



# Cloudy Results: Ascending SST



- Results are similar for v5.4 & v5.4Clim
- Missed clouds are the main problem for both
  - V5.4 is more prone to run away >20K cold
    - These would normally be identified by QC
  - V5.4Clim sometimes just fails in cloud clearing for these runaway cases because the startup state is too far from the truth






National Aeronautics and  
Space Administration  
Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

# Cloudy Results: Ascending SST

*Atmospheric Infrared Sounder*

v5.4

SST - Truth



5/4/09

Copyright 2009 California Institute of Technology.  
Government sponsorship acknowledged.

15






National Aeronautics and  
Space Administration  
Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, California

# Cloudy Results: Ascending SST

*Atmospheric Infrared Sounder*

v5.4 Clim

SST - Truth




5/4/09

Copyright 2009 California Institute of Technology.  
Government sponsorship acknowledged.

16




# Cloudy Results: Temperature Profile Skill



- Global stats suggest v5.4Clim is superior for skill, bias, and standard deviation of temperature profile throughout the troposphere.
- The difference is greatest at the surface
  - V5.4Clim seems unaffected by problems with cloud clearing

# Cloudy Results: Temperature Profile Skill



- V5.4Clim seems much better than v5.4 in the tropics (+/- 20 degrees latitude) where the variation of the weather is relatively small and cloudiness is high.
- V5.4Clim seems best below 700 mbar everywhere.
- V5.4 is better in the troposphere above 700 mbar in the moderate region (40-60 degrees latitude) where the weather is more changeable
- Results for the subtropical and polar zones also favor v5.4Clim, though not as strongly as the tropics do.

# Analysis: Climatology Startup

- V5.4Clim compares surprisingly well with the v5.4 baseline, even without changes to the physical retrieval
- A well-tuned system with a climatology start-up would need:
  - Substantially more channels than the baseline system
    - Because regression is currently providing all vertical fine structure
  - Substantially more vertical retrieval levels than the baseline system
    - Because the initial state is further from the final state
  - Substantially more CPU time
    - This slowdown is acceptable.

# Analysis: Extreme Weather

- Extreme weather conditions are the most difficult for a system with a climatology startup
  - Cloud clearing must work reasonably with a background state far from the true state
  - The physical retrieval must make large changes
- The skill metric is specifically designed to weight non-typical weather patterns more highly
- The v5.4Clim climatology startup system already does well, even in terms of skill, compared to the v5.4 regression startup system
  - Apparently extreme weather conditions are also a problem for regression

# SCCNN

- Bill Blackwell has described the algorithm at previous meetings.  
[http://airs.jpl.nasa.gov/documents/science\\_team\\_meeting\\_archive/science\\_team\\_meeting\\_2008.04.15/science\\_team\\_meeting\\_2008.04.15\\_files/blackwell.pdf](http://airs.jpl.nasa.gov/documents/science_team_meeting_archive/science_team_meeting_2008.04.15/science_team_meeting_2008.04.15_files/blackwell.pdf)
  - Regression-like
- Last year I showed results of skill testing on SCCNN
  - Results were better than regression and almost equal to full retrieval
  - The boundary layer was the main problem area
  - Newer test data shows the boundary layer problem is overcome
- MIT has produced a FORTRAN-language version of the algorithm suitable for integration in AIRS operational Level-2 retrieval code
- Jan Gohlke has the standalone program running at JPL in preparation for integration.
  - Final debugging and validation is still in process.

# Continuing Work

- Climatology needs to be completed.
  - Currently have only 6 months from 1-2 years
    - Need all 12 months
    - Need to fill in gaps in recent ECMWF forecast
  - Evaluate use of multiyear average vs. latest
- The physical retrieval must be optimized to work with climatology startup before a full comparison can be made:
  - Add channels and slabs
  - Reduce damping/increase iterations
  - Use climate variance as an error estimate on startup  $T(p)$ ,  $q(p)$
- If regressions are going to continue to be used their training set needs to be updated to include more recent data
- We need to compare baseline v6 with:
  - SCCNN
  - Climatology
    - Replicate results at other locations
  - MW-Only with the new climatology

# Conclusions


- Climatology startup is a viable alternative to regression.
  - More development and testing is needed
- The final choice of v6 startup should not be based only on skill or other measurements of accuracy and yield. We also must consider:
  - Predictability of results
  - Error propagation
  - Availability of averaging kernels

# BACKUP




# SST v5.4 Clim – Climatology: Shows Physical Retrieval Can Make Large Changes

2008.12.07.L3.RetStd\_IR008:SurfSkinTemp\_A – 2007.12.01.L3.RetStd031:SurfSkinTemp\_A



Difference Range: -23.9766 to 26.6875



5/4/09

Copyright 2009 California Institute of Technology.  
 Government sponsorship acknowledged.